

QUANTUM Z -ALGEBRAS AND REPRESENTATIONS OF QUANTUM AFFINE ALGEBRAS

NAIHUAN JING

Department of Mathematics
North Carolina State University
Raleigh, NC 27695-8205, U.S.A.

ABSTRACT. Generalizing our earlier work, we introduce the homogeneous quantum Z -algebras for all quantum affine algebras $U_q(\widehat{\mathfrak{g}})$ of type one. With the new algebras we unite previously scattered realizations of quantum affine algebras in various cases. As a result we find a realization of $U_q(F_4^{(1)})$.

0. INTRODUCTION

In 1981 Lepowsky and Wilson introduced (principal) Z -algebras as a tool to construct explicit principal representations of affine Lie algebras [LW1]. Later Lepowsky and Primc considered the homogeneous analog for $\widehat{sl}_2$ and gave explicit realizations of higher level representations of $\widehat{sl}_2$ [LP1, LP2]. Such a construction is also done for the higher rank cases [P]. Many other vertex representations of affine algebras were constructed by this method (cf. [FLM] for a list of references).

Recently we introduced the quantum Z -algebras for the simply laced types in the homogeneous picture [J2] and used them to construct higher level (integral) representations of the quantum affine algebra $U_q(\widehat{sl}_2)$. In [BV] the higher level representations of $U_q(\widehat{sl}_2)$ and its Z -algebra in the Verma module picture were also studied. These constructions parallel to the classical method of Lepowsky and Primc. As we mentioned in [J2], the quantum Z -algebras are expected to provide more examples of explicit constructions of quantum affine algebras.

In this paper we define the quantum Z -algebras for all type one quantum affine algebras. The quantum Z -algebras enable us to unify previously scattered constructions of quantum affine algebras, and we also find a realization of $U_q(F_4^{(1)})$ based on our recent joint work on representations of $U_q(C_n^{(1)})$ [JKM1, JKM2]. The latter also contained some quantum deformation of the admissible representations studied by Kac and Wakimoto in the classical case (in this special case, see Feingold-Frenkel [FF]). Thus all level one representations of quantum affine algebras of type one can be constructed by vertex operators and quantum Z -algebras. Since we work in a general setting we can derive all commutation relations for the quantum Z -algebras at once with the help of q -series. As an simple example we recover the Frenkel-Jing

1991 *Mathematics Subject Classification.* 17B.

Key words and phrases. Quantum enveloping algebras, Quantum Z -operators, Vertex operators, Hopf algebras.

bosonic realizations of quantum affine algebras of simply laced types. As for other cases we try to emphasize on the similarity among all the constructions and points out differences in each special case.

This paper mainly concerns with homogeneous quantum Z -algebras. The quantum principal Z -algebra is still unknown, though the related fermionic constructions are known quite generally (cf. [KMPY]), where one can also find more examples about higher level $U_q(\widehat{sl}_2)$ -modules.

1. QUANTUM AFFINE ALGEBRAS AND THEIR MODULES

Let $\widehat{\mathfrak{g}}$ be the affine Kac-Moody algebra [K] associated to generalized Cartan matrix of type one $A = (A_{ij})$. Let $\mathfrak{g}$ be the associated finite dimensional simple Lie algebra. Let $Q = \mathbb{Z}\alpha_1 \oplus \cdots \oplus \mathbb{Z}\alpha_l$ and $P = \mathbb{Z}\lambda_1 \oplus \cdots \oplus \mathbb{Z}\lambda_l$ be the root and weight lattices of $\mathfrak{g}$. We fix the standard normalized invariant $(\mid)$ on $\mathfrak{g}$ such that

$$(\alpha_i \mid \alpha_j) = d_i A_{ij}, \quad i, j = 1, \dots, l$$

where the $d_i = (\alpha_i \mid \alpha_i)/2$. Let $q_i = q^{d_i}$.

The quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ is the associative algebra generated by the central element γ , the grading element d , and the elements $\alpha_i(n)$, and $x_i^\pm(n)$, $i = 1, \dots, l$ (or equivalently the elements $x^\pm(n)$, $\phi(-m)$, $\psi(m)$), $n \in \mathbb{Z}$ and $m \in \mathbb{Z}_+$ subject to the following relations:

$$\begin{aligned} [d, a_i(n)] &= n\alpha_i(n), [d, x_i^\pm(n)] = nx_i^\pm(n), \\ [a_i(m), a_j(n)] &= \delta_{m,-n} \frac{[(\alpha_i \mid \alpha_j)m]}{m} \frac{\gamma^m - \gamma^{-m}}{q - q^{-1}}, \\ [a_i(m), x_j^\pm(n)] &= \pm \frac{[(\alpha_i \mid \alpha_j)m]}{m} \gamma^{\mp |m|/2} x_j^\pm(m+n), \\ (z - q^{\pm A_{ij}} w) x_i^\pm(z) x_j^\pm(w) &= x_j^\pm(w) x_i^\pm(z) (q^{\pm A_{ij}} z - w) \\ [x_i^+(z), x_j^-(w)] &= \frac{\delta_{ij}}{q_i - q_i^{-1}} \left\{ \delta\left(\frac{z}{w} q^{-c}\right) \psi_i(w q^{c/2}) - \delta\left(\frac{z}{w} q^c\right) \phi_i(z q^{c/2}) \right\}, \\ Sym_{z_1, \dots, z_N} \sum_{s=0}^{N=1-A_{ij}} (-1)^s \begin{bmatrix} N \\ s \end{bmatrix}_i x_i^\pm(z_1) \cdots x_i^\pm(z_s) \cdot \\ &\cdot x_j^\pm(w) x_i^\pm(z_{s+1}) \cdots x_i^\pm(z_N) = 0, \quad \text{for } A_{ij} \neq 0 \end{aligned}$$

where the generators $\alpha(n)$ are related to $\phi(-m)$, $\psi(m)$ via:

$$\begin{aligned} \phi_i(z) &= \sum_{n \geq 0} \phi_i(-n) z^n = k_i^{-1} \exp((q^{-1} - q) \sum_{n > 0} \alpha_i(-n) z^n), \\ \psi_i(z) &= \sum_{n \geq 0} \psi_i(n) z^{-n} = k_i \exp((q - q^{-1}) \sum_{n > 0} \alpha_i(n) z^{-n}). \end{aligned}$$

The fields are defined by

$$x_i^\pm(z) = \sum x_i^\pm(n) z^{-n-1}, \quad i = 1, \dots, l.$$

We have rescaled the generators $a_i(n)$ by the factor $[(\alpha_i|\alpha_i)/2]$ to have an invariant presentation. With $a_i(n)/[(\alpha_i|\alpha_i)/2]$ as $a_i(n)$ we get the Drinfeld realization studied in [Be, J4].

We define the q -integers by

$$[n]_i = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}}, \quad \left[\begin{matrix} m \\ n \end{matrix} \right]_i = \frac{[m]_i!}{[n]_i! [m-n]_i!}.$$

We also need the q -number (for $a \in \mathbb{R}$):

$$(a; q)_n = (1-a)(1-aq) \cdots (1-q^{n-1}a).$$

Let V be a $U_q(\widehat{\mathfrak{g}})$ -module. The module V is called level k if the central element γ acts as the scalar q^k on V . For $a \in \mathbb{C}$, define

$$V_a = \{v \in V | d.v = av\},$$

the elements of V_a are homogeneous elements of degree a . An endomorphism A of V is homogeneous of degree $n \in \mathbb{C}$ if

$$[d, A] = nA.$$

We will study the following category of $U_q(\widehat{\mathfrak{g}})$ -modules V with the properties:

- (1) the module V has level k ;
- (2) each weight space of V is finite dimensional;
- (3) for each $a \in \mathbb{C}$, there exists $n_0 \geq 0$ such that for all $n > n_0$, $V_{a+n} = \{0\}$.

It is clear that the irreducible highest weight modules $L(\lambda)$ lie in the category for a suitable k . The level is an integer for the dominant integrable module.

2. QUANTUM Z-ALGEBRAS

Let V be a $U_q(\widehat{\mathfrak{g}})$ -module of level k , $k \in \mathbb{Q}$. The Heisenberg algebra $U_q(\widehat{\mathfrak{h}})$ of level k is generated by $\alpha_i(n)$, $n \neq 0$ and the central element c subject to the relations:

$$(2.1) \quad [\alpha_i(m), \alpha_j(n)] = \frac{1}{m} [(\alpha_i|\alpha_j)m][mk]\delta_{m,-n}.$$

and V is also a $U_q(\widehat{\mathfrak{h}})$ -module.

We define the following operators on V :

$$(2.2) \quad \begin{aligned} E_-^\pm(\alpha_i, z) &= \exp(\mp \sum_{n \geq 1} \frac{q^{\mp kn/2}}{[kn]} \alpha_i(-n) z^n) \\ E_+^\pm(\alpha_i, z) &= \exp(\pm \sum_{n \geq 1} \frac{q^{\mp kn/2}}{[kn]} \alpha_i(n) z^{-n}) \end{aligned}$$

For $a \in \mathbb{R}$ we set

$$(1-z)^{a/k}_{q^{2k}} = \frac{(q^{-a+k}z; q^{2k})_\infty}{(q^{a+k}z; q^{2k})_\infty} = \exp(-\sum_{n=1}^{\infty} \frac{[an]}{n[kn]} z^n)$$

The Heisenberg algebra has a natural realization on the space $Sym(\widehat{\mathfrak{h}}^-)$ of the symmetric algebra generated by the elements $\alpha_i(-m)$, $m \in \mathbb{N}$ via the following rule:

$$\begin{aligned} q^c \cdot 1 &= q^k, & q^{-c} \cdot 1 &= q^{-k}, \\ \alpha_i(-n) &= \text{multiplication operator}, & n &\in \mathbb{N} \\ \alpha_i(n) &= \text{differentiation operator subject to (2.1)}. \end{aligned}$$

We denote the resulted irreducible module of $U_q(\widehat{\mathfrak{h}})$ as $K(k)$.

We now introduce the Z -vertex operators. Let V be a level k $U_q(\widehat{\mathfrak{g}})$ -module. For each $i = 1, \dots, l$, define

$$(2.5) \quad Z^\pm(\alpha_i, z) = E_-^\pm(\alpha_i, z) x_i^\pm(z) E_+^\pm(\alpha_i, z),$$

which act on the space V .

Proposition 2.1. *As operators on a level k module V ,*

$$\begin{aligned} E_+^\pm(\alpha_i, z) E_-^\pm(\alpha_j, w) &= E_-^\pm(\alpha_j, w) E_+^\pm(\alpha_i, z) \left(1 - \frac{w}{z} q^{\mp k}\right)_{q^{2k}}^{(\alpha_i|\alpha_j)/k}, \\ E_+^\pm(\alpha_i, z) E_-^\mp(\alpha_j, w) &= E_-^\mp(\alpha_j, w) E_+^\pm(\alpha_i, z) \left(1 - \frac{w}{z}\right)_{q^{2k}}^{-(\alpha_i|\alpha_j)/k}, \\ E_+^\pm(\alpha_i, z) x_j^\pm(w) &= x_j^\pm(w) E_+^\pm(\alpha_i, z) \left(1 - \frac{w}{z} q^{\mp k}\right)_{q^{2k}}^{-(\alpha_i|\alpha_j)/k}, \\ E_+^\pm(\alpha_i, z) x_j^\mp(w) &= x_j^\mp(w) E_-^\pm(\alpha_i, z) \left(1 - \frac{w}{z}\right)_{q^{2k}}^{(\alpha_i|\alpha_j)/k}, \\ E_+^\pm(\alpha_i, z) x_j^\pm(w) &= x_j^\pm(w) E_+^\pm(\alpha_i, z) \left(1 - \frac{w}{z} q^{\mp k}\right)^{-(\alpha_i|\alpha_j)/k}, \\ [\phi(\alpha_i, z), E_-^\pm(\alpha_j, w)] &= [\psi(\alpha_i, z), E_+^\pm(\alpha_j, w)] = 0, \\ E_+^\pm(\alpha_j, w) \phi(\alpha_i, z) &= \phi(\alpha_i, z) E_+^\pm(\alpha_j, w) \frac{w - q^{\pm(\alpha_i|\alpha_j) \mp k/2} z}{w - q^{\mp(\alpha_i|\alpha_j) \mp k/2} z}, \\ \psi(\alpha_i, z) E_-^\pm(\alpha_j, w) &= E_-^\pm(\alpha_j, w) \psi(\alpha_i, z) \frac{z - q^{\pm(\alpha_i|\alpha_j) \mp k/2} w}{z - q^{\mp(\alpha_i|\alpha_j) \mp k/2} w}, \\ E_-^+(\alpha_i, z) E_-^-(\alpha_i, z q^{-k}) &= 1, \\ E_-^+(-\alpha_i, z) E_-^-(\alpha_i, z q^k) &= \phi(\alpha_i, z q^{k/2}) k_i, \\ E_+^+(-\alpha_i, z) E_+^-(\alpha_i, z q^{-k}) &= \psi(\alpha_i, z q^{-k/2}) k_i^{-1}, \\ E_+^+(\alpha_i, z) E_+^-(\alpha_i, z q^k) &= 1. \end{aligned}$$

□

Theorem 2.2. *As operators on a level k space, the Z -vertex operators satisfy the following relations.*

$$\begin{aligned} &(1 - q^{\mp k} w/z)_{q^{2k}}^{-(\alpha_i|\alpha_j)/k} (z - q^{\pm(\alpha_i|\alpha_j)} w) Z^\pm(\alpha_i, z) Z^\pm(\alpha_j, w) \\ &= (1 - q^{\mp k} z/w)_{q^{2k}}^{-(\alpha_i|\alpha_j)/k} (q^{\pm(\alpha_i|\alpha_j)} z - w) Z^\pm(\alpha_j, w) Z^\pm(\alpha_i, z), \\ &(1 - w/z)_{q^{2k}}^{(\alpha_i|\alpha_j)/k} Z^+(\alpha_i, z) Z^-(\alpha_j, w) - (1 - z/w)_{q^{2k}}^{(\alpha_i|\alpha_j)/k} Z^-(\alpha_j, w) Z^+(\alpha_i, z) \\ &= \frac{\delta_{ij}}{-1} \{k_i \delta(q^{-k} z/w) - k_i^{-1} \delta(q^k z/w)\} \end{aligned}$$

Proof. The relations are consequences of the relations satisfied by the operators $E_{-}^{\pm}(z)$ and $E_{+}^{\pm}(z)$. For details, see [J2] $\square$

We will call the algebra generated by operators satisfying the relations of Theorem 2.2 as the quantum Z -algebra. We may also require that the relations contain the Serre relations (which are derived from the Serre relations of $X_i^{\pm}(z)$) to ensure that it will come from a representation of the quantum affine algebra. We define the vacuum space Ω_V as an invariant subspace of the quantum Z -operators.

$$\Omega_V = \{v \in V \mid E_{+}^{\pm}(\alpha_i, z).v = E_{+}^{\pm}(-\alpha_i, z).v = v\}.$$

As in the simply laced case [J2] we have

Theorem 2.3. *Let V be a level k module, then we have a linear isomorphism:*

$$\begin{aligned} K(k) \otimes_{\mathbb{C}} \Omega_V &\longrightarrow V \\ u \otimes w &\longmapsto i(u) \cdot w \end{aligned}$$

where we identify the space $K(k)$ with the Heisenberg algebra via the classical action, and the inclusion i embeds $U_q(\widehat{\mathfrak{h}}^-)$ into $U_q(\widehat{\mathfrak{g}})$. Subsequently we can write

$$x_i^{\pm}(z) = E_{-}^{\pm}(-\alpha_i, z)E_{+}^{\pm}(-\alpha_i, z) \otimes Z^{\pm}(\alpha_i, z).$$

Proof. The proof is exactly the same as in the simply laced case, as we observe that $Z^{\pm}(\alpha_i, z)$ commutes with the operators $a_i(n)$ of the Heisenberg subalgebra. $\square$

Thus the construction of irreducible modules is reduced to that of corresponding Z -vertex operators.

Proposition 2.4. *A $U_q(\widehat{\mathfrak{g}})$ -module V is irreducible if and only if the vacuum space Ω_V is an irreducible $U_q(\widehat{\mathfrak{h}})$ -module.* $\square$

3. VARIOUS CONSTRUCTIONS OF $U_q(\widehat{\mathfrak{g}})$

3.1. $k = 1$, Frenkel-Jing realization for $(ADE)^{(1)}$ [FJ].

Let $\mathbb{C}\{Q\}$ be the twisted group algebra of Q generated by e^{α} , $\alpha \in Q$ subject to the relations

$$e^{\alpha}e^{\beta} = (-1)^{(\alpha|\beta)}e^{\beta}e^{\alpha}.$$

For each level one integrable module Λ_r , there corresponds to a minuscule weight $\lambda_r \in Q$ (cf. Table 1.1). We formally adjoin e^{λ_r} to form the vector space $\mathbb{C}\{Q\}e^{\lambda_r}$.

We define the action of $z^{\partial_{\alpha}}$ on $\mathbb{C}\{Q\}e^{\lambda_r}$ by

$$z^{\partial_{\alpha}}e^{\beta}e^{\lambda_r} = z^{(\alpha|\beta+\omega_r)}e^{\beta}e^{\lambda_r}$$

for $\alpha \in Q$.

The following construction is a reformulation of the basic vertex representations given in [J2].

The level one module $V(\Lambda_r)$ is realized as

$$V = \text{Sym}(\widehat{\mathfrak{h}}^-) \otimes \mathbb{C}\{Q\}e^{\lambda_r}$$

with the vacuum space equals to $\mathbb{C}\{Q\}e^{\lambda_r}$ and the Z -operators are given as:

$$Z^{\pm}(\alpha_i) = e^{\pm\alpha_i}z^{\pm\partial_{\alpha_i}},$$

and the highest weight vector is e^{λ_r} .

The construction was generalized to all twisted types in [J1]

3.2. $B_l^{(1)}$, $k=1$, [B, JM].

There are three level one modules: $V(\Lambda_0)$, $V(\Lambda_1)$, and $V(\Lambda_l)$. We can write $\Lambda_i = \Lambda_0 + \lambda_i$, where λ_i are the fundamental weights for $\mathfrak{g}$.

Let $\mathbb{C}\{Q\}$ be the twisted group algebra generated by e^α , $\alpha \in Q$ subject to the relations:

$$e^\alpha e^\beta = (-1)^{(\alpha|\beta) + (\alpha|\alpha)(\beta|\beta)} e^\beta e^\alpha.$$

The vector space $\mathbb{C}\{P\}$ is defined as in §3.1. Let Q_0 be the sublattice of the long roots, then we have

$$\mathbb{C}\{Q\} = \mathbb{C}\{Q_0\} \oplus \mathbb{C}\{Q_0\}e^{\lambda_1}.$$

We set $Z = \mathbb{Z} + s$, $s = 0$ or $1/2$. The q -deformed Clifford algebra C_q^s is the quadratic algebra generated by $\kappa(k)$ satisfying the relations:

$$(3.1) \quad \{\kappa(m), \kappa(n)\} = (q^m + q^{-m})\delta_{m,-n},$$

where $\{a, b\}$ is the anticommutator, and $m, n \in Z$.

Let $\Lambda(C_q^-)^s$ be the polynomial algebra generated by $\kappa(-k)$, $k \in Z_{>0}$, and $\Lambda(C_q^-)_0^s$ (resp. $\Lambda(C_q^-)_1^s$) be the subalgebra generated by products of even (resp. odd) number of generators $\kappa(-k)$'s. Then

$$\Lambda(C_q^-)^s = \Lambda(C_q^-)_0^s \oplus \Lambda(C_q^-)_1^s.$$

The algebra C_q^s acts on $\Lambda(C_q^-)^s$ canonically by ($k \in Z_{>0}$)

$$\begin{aligned} \kappa(-n).v &= \kappa(-n) \wedge v, & \text{for } n > 0 \\ \kappa(n).1 &= 0 & \text{for } n > 0 \end{aligned}$$

We define that

$$\begin{aligned} W_0 &= \text{Sym}(\hat{\mathfrak{h}}^-) \otimes (\mathbb{C}\{Q_0\} \otimes \Lambda(C_q^-)_0^{1/2} \oplus \mathbb{C}\{Q_0\}e^{\lambda_1} \otimes \Lambda(C_q^-)_1^{1/2}), \\ W_1 &= \text{Sym}(\hat{\mathfrak{h}}^-) \otimes (\mathbb{C}\{Q_0\}e^{\lambda_1} \otimes \Lambda(C_q^-)_0^{1/2} \oplus \mathbb{C}\{Q_0\} \otimes \Lambda(C_q^-)_1^{1/2}), \\ W_l &= \text{Sym}(\hat{\mathfrak{h}}^-) \otimes \mathbb{C}\{Q\}e^{\lambda_l} \otimes \Lambda(C_q^-)^0. \end{aligned}$$

Then the algebras $U_q(\hat{\mathfrak{h}})$, $\mathbb{C}\{P\}$ and C_q^s (for suitable $s = 0, 1/2$) act on the spaces W_i canonically by extending their respective actions component-wise. We also define the degree action by

$$d.f \otimes e^\alpha \otimes g = \left(-\sum_{j=1}^k m_j - \sum_{j=1}^l n_j - \frac{(\beta|\beta)}{2} + \frac{(\alpha_i|\alpha_i)}{2}\right) f \otimes e^\alpha \otimes g,$$

where $f \otimes e^\alpha \otimes g = a_{i_1}(-m_1) \cdots a_{i_k}(-m_k) \otimes e^\alpha \otimes \kappa(-n_1) \cdots \kappa(-n_l) \in W_i$. For convenience we denote $\lambda_0 = 0$.

Proposition 3.2. *The space W_i ($i = 0, 1, l$) is the irreducible representation $V(\Lambda_i)$ of the quantum affine Lie algebra $U_q(B_l^{(1)})$ under the action:*

$$\begin{aligned} \gamma &\mapsto q, \quad K_j \mapsto q^{\partial_{\alpha_j}}, \quad a_j(k) \mapsto a_j(k) \quad (1 \leq j \leq l), \\ Z_i^\pm(z) &= e^{\pm\alpha_i} z^{\pm\partial_i}, \quad i = 1, \dots, l-1 \\ Z_l^\pm(z) &= e^{\pm\alpha_l} z^{\pm\partial_l} (\pm\kappa(z)), \end{aligned}$$

where $\kappa(z) = \sum_{m \in Z} \kappa(m)z^{-m}$ and the degree operator d is defined above. The highest weight vectors are respectively:

$$|\Lambda_i\rangle = 1 \otimes e^{\lambda_i} \otimes 1, \quad i = 0, 1, l.$$

3.3. $k = 2$, $A_1^{(1)}$ [I, J2]

There are three level two integrable modules $V(2\Lambda_0)$, $V(2\Lambda_1)$, and $V(2\Lambda_0 + 2\Lambda_1)$.

We need the Clifford algebra $C_{q^2}^{1/2}$ (3.1) to realize level 2 $U_q(\widehat{sl}_2)$.

Define the action of d on $\Gamma(\mathcal{C}_{q^2}^s)$ by

$$d.\kappa(n_1) \wedge \cdots \wedge \kappa(n_m) = (n_1 + \cdots + n_m)\kappa(n_1) \wedge \cdots \wedge \kappa(n_m).$$

Proposition 3.3. [J2] *Set*

$$\Omega = \Gamma(\mathcal{C}_{q^2}^{-,1/2}) \otimes \mathbb{C}[Q],$$

and extend naturally the actions of e^β and d to the space Ω . We can define a representation of $U_q(\widehat{sl}_2)$ on $V = K(2) \otimes \Omega$ via:

$$Z^\pm(z) = \pm\kappa(z) \otimes e^{\pm\alpha} z^{\pm\alpha/2+1/2}.$$

Moreover, the modules V decomposes itself as

$$V = L(2\Lambda_0) \oplus L(2\Lambda_1 - (1/2)\delta).$$

For the level two module $L(\Lambda_0 + \Lambda_1)$ we consider another Clifford algebra $\mathcal{C}_{q^2}^0$.

In order to construct the vacuum space for the representation $L(\Lambda_0 + \Lambda_1)$ we need to extend the space $\Gamma(\mathcal{C}_{q^2}^{-,0})$. Define the action of $\kappa(0)$ on the two-dimensional space $\mathbb{C}^2$ by

$$\kappa(0) \cdot \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} v \\ u \end{pmatrix}$$

and then naturally extend the action of $\mathcal{C}$ on the space $\Gamma(\mathcal{C}_{q^2}^{-,0}) \otimes \mathbb{C}^2$.

Proposition 3.4. [J2] *Let $V = K(2) \otimes \Omega_V$ with*

$$\Omega_V = \Gamma(\mathcal{C}_{q^2}^{-,0}) \otimes \mathbb{C}^2 \otimes \mathbb{C}[Q]e^{\alpha/2},$$

then we can realize a representation of $U_q(\widehat{sl}_2)$ by

$$Z^\pm(z) = \pm\kappa(z) \otimes e^{\pm\alpha} z^{\pm\alpha/2+1/2}.$$

Moreover, we have

$$V = L(\Lambda_0 + \Lambda_1 - \delta/8) \oplus L(\Lambda_0 + \Lambda_1 - \delta/8).$$

Proof. The proposition is proved similarly as Proposition 3.2. □

3.4. Higher level $U_q(\widehat{sl}_2)$ [J2, ABH, AOS, KMPY, M, S]

We consider irreducible modules of highest weights

$$\lambda = i\Lambda_0 + (k - i)\Lambda_1,$$

where $i \in \{0, \dots, k\}$. We construct the module as an submodule of the tensor product of level one modules

$$(3.2) \quad L(\Lambda_0)^{\otimes i} \otimes L(\Lambda_1)^{\otimes (k-i)}$$

where the tensor product is induced from the Drinfeld comultiplication (cf. [DFr] [J2]). Drinfeld's coproduct plays an important role in studying higher level representations based on Frenkel-Jing constructions as we noticed in [J2]. This technique was further developed by the work [DW, DFg].

In order to state Drinfeld comultiplication we introduce the following notation. For a field operator $A(z) = \sum_n A_n z^{-n}$ acting on a space V and a commutative operator q^c on V we define

$$\begin{aligned} A(z \otimes q^c) &= \sum_n (A_n \otimes q^{nc}) z^{-n} = \sum_n A_n z^{-n} \otimes q^{nc} \\ A(q^c \otimes z) &= \sum_n (q^{nc} \otimes A_n) z^{-n} = \sum_n q^{nc} \otimes A_n z^{-n} \end{aligned}$$

as operators on $V \otimes V$. In particular, $A(z \otimes 1) = A(z) \otimes 1$. The Drinfeld comultiplication is given by the following formula:

$$(3.3) \quad \Delta(\phi_i(z)) = \phi_i(z \otimes q^{-c/2}) \phi_i(q^{c/2} \otimes z),$$

$$(3.4) \quad \Delta(\psi_i(z)) = \psi_i(z \otimes q^{c/2}) \psi_i(q^{-c/2} \otimes z),$$

$$(3.5) \quad \Delta(x_i^+(z)) = x_i^+(z \otimes 1) + \psi_i(q^{-c/2} z \otimes 1) x_i^+(q^{-c} \otimes z),$$

$$(3.6) \quad \Delta(x_i^-(z)) = x_i^-(1 \otimes z) + x_i^-(z \otimes q^{-c}) \phi_i(1 \otimes q^{-c/2} z).$$

Thus the first formula means that

$$\begin{aligned} \Delta(\phi_i(z)) &= \sum_{m,n} (\phi_i(-n) z^n \otimes q^{-nc/2}) \cdot (q^{mc/2} \otimes \phi_i(-m) z^m) \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^n \phi_i(-m) q^{(n-m)c/2} \otimes \phi_i(-n+m) q^{-mc/2} z^n. \end{aligned}$$

The counit ϵ and antipode S are given by

$$\begin{aligned} \epsilon(1) &= \epsilon(\phi_i(z)) = \epsilon(\psi_i(z)) = 1, \\ \epsilon(x_i^{\pm}(z)) &= 0, \\ S(\phi_i(z)) &= \phi_i(z)^{-1}, \quad S(\psi_i(z)) = \psi_i(z)^{-1}, \\ S(x_i^+(z)) &= -\psi_i(q^{c/2} z)^{-1} x_i^+(q^c z) \\ S(x_i^-(z)) &= -x_i^-(q^{-c} z)^{-1} \phi_i(q^{c/2} z)^{-1} \end{aligned}$$

In [J2] we showed that the comultiplication is equivalent to the following component form:

$$(3.7) \quad \Delta(\alpha_i(n)) = q^{|n|c/2} \otimes \alpha_i(n) + \alpha_i(n) \otimes q^{-|n|c/2},$$

$$(3.8) \quad \Delta(x_i^+(n)) = x_i^+(n) \otimes 1 + \sum_{m \geq 0} \psi_i(m) q^{(m+2n)c/2} \otimes x_i^+(n-m),$$

$$(3.9) \quad \Delta(x_i^-(n)) = 1 \otimes x_i^-(n) + \sum_{m \geq 0} x_i^-(n+m) \otimes \phi_i(-m) q^{(m+2n)c/2}.$$

We extend the comultiplication to multi-tensor product in the natural way. Define

$$\Delta_{k-1} : U_q(\widehat{\mathfrak{g}}) \longrightarrow U_q(\widehat{\mathfrak{g}})^{\otimes k}$$

inductively

$$\Delta_1 = \Delta, \Delta_j = (1 \otimes \Delta) \Delta_{j-1} = (\Delta \otimes 1) \Delta_{j-1}, \quad \text{for } j = 2, \dots, k-1$$

For $\epsilon_1, \dots, \epsilon_s \in \{\pm\} = \{\pm 1\}$ we have

$$\begin{aligned} X_1^{\epsilon_1}(z_1) \cdots X_s^{\epsilon_s}(z_s) &= E_-^{\epsilon_1}(-\alpha_1, z_1) \cdots E_-^{\epsilon_s}(-\alpha_s, z_s) \\ &\quad E_+^{\epsilon_1}(-\alpha_1, z_1) \cdots E_+^{\epsilon_s}(-\alpha_s, z_s) Z(\epsilon_1 \cdots \epsilon_s; z_1 \cdots z_s) \end{aligned}$$

where the-multi argument Z -operator $Z(\epsilon_1 \cdots \epsilon_s; z_1 \cdots z_s)$ is defined by

$$Z(\epsilon_1 \cdots \epsilon_s; z_1 \cdots z_s) = \prod_{1 \leq i < j \leq s} \left(1 - \frac{z_j}{z_i} q^{-(\epsilon_i + \epsilon_j)k/2}\right)^{-2\epsilon_i \epsilon_j / k} Z^{\epsilon_1}(\alpha_1, z_1) \cdots Z^{\epsilon_s}(\alpha_s, z_s).$$

In particular, we have

$$Z(\epsilon, z) = Z^\epsilon(\alpha, z).$$

We define the operators $Z(\epsilon_1 \cdots \epsilon_s; n_1 \cdots n_s)$ as the components of the Z -operators:

$$Z(\epsilon_1 \cdots \epsilon_s; z_1 \cdots z_s) = \sum Z(\epsilon_1 \cdots \epsilon_s; n_1 \cdots n_s) z^{-n_1} \cdots z^{-n_s}.$$

Clearly the operators $Z(\epsilon_1 \cdots \epsilon_s; n_1 \cdots n_s)$ preserve Ω_V , and

$$\deg Z(\epsilon_1 \cdots \epsilon_s; n_1 \cdots n_s) = n_1 + \cdots + n_s.$$

Using the results of poles [DW] and Drinfeld comultiplication, Ding and Feigin [DFg] recently gave a comprehensive study of higher level representations of the quantum affine algebra $U_q(\widehat{sl_2})$.

In the above we consider examples of irreducible modules. We now turn to reducible modules. According to Proposition 2.4 we only need to construct quantum Z -algebras acting on $U_q(\widehat{\mathfrak{h}})$ -reducible modules. Usually we need to enlarge the algebra by adding auxiliary operators. The quantum Wakimoto constructions are done in this way. In the following we will only discuss one of the constructions, since all are closely related [Bo]. Level k modules of $\widehat{sl_2}$ are also studied in [KMPY] using crystal graphs.

Consider three independent sets of bosonic operators $a(n), b(n), c(n)$ satisfying the commutation relations:

$$\begin{aligned} [a(n), a(m)] &= \frac{[2n]}{n} [nk] \delta_{n,-m} \\ [b(n), b(m)] &= -\frac{[2n]}{n} [nk] \delta_{n,-m} \\ [c(n), c(m)] &= \frac{[2n]}{n} [n(k+2)] \delta_{n,-m} \end{aligned}$$

and we let β be an auxiliary root such that $(\beta|\beta) = -(\alpha|\alpha)$ and $(\alpha|\beta) = 0$.

Proposition 3.5. *The realization space of $U_q(A_1^{(1)})$ at level k is defined as*

$$V = \text{Sym}(a(-n)'s, b(-n)'s, c(-n)'s) \otimes C[\mathbb{Z}\alpha + \mathbb{Z}\beta].$$

and the action is given by

$$\begin{aligned} Z^\pm(z) \mapsto e^{\pm\alpha_1} z^{\pm\partial_\alpha} \frac{U^\pm(z)}{q - q^{-1}} & (B_+(zq^{\mp(k+2)/2}z)W_+(q^{\mp k/2}z)^{\pm 1} \\ & - B_-(zq^{\pm(k+2)/2}z)W_-(q^{\pm k/2}z)^{\pm 1}), \end{aligned}$$

where

$$\begin{aligned} U^\pm(z) &= \exp(\pm \sum_{m=1}^{\infty} \frac{q^{\mp km/2} b(-m)}{[km]} z^m) \exp(\mp \sum_{m=1}^{\infty} \frac{q^{\mp km/2} b(m)}{[km]} z^{-m}) e^{\pm\beta} z^{\pm\partial_\beta} \\ B_\pm(z) &= \exp(\pm(q - q^{-1}) \sum_{m=1}^{\infty} \frac{[m]}{[2m]} b(\pm) z^{\mp m}) q^{\mp\partial_\beta} \\ W_\pm(z) &= \exp(\pm(q - q^{-1}) \sum_{m=1}^{\infty} \frac{[m]}{[2m]} c(\pm) z^{\mp m}). \end{aligned}$$

3.5. Realizations of $U_q(C_l^{(1)})$ and $U_q(F_4^{(1)})$.

Recently we have constructed the quantum symplectic affine algebra $U_q(C_n^{(1)})$ at level $-1/2$ [JKM1] and level 1 [JKM2]. The reducible modules were constructed using the same idea of the quantum Z -algebras. In the case of level one our realization contains all integrable level one modules.

Recalling the construction in 3.1 we also formally define the space $C[P]$ as the direct sum $C[Q] \oplus C[Q]e^{\lambda_1}$ by adjoining the element e^{λ_1} . The short roots Q_s constitute the root system A_{n-1} . Let $\tilde{Q}$ be the root lattice of A_{l-1} generated $\tilde{\alpha}_i$, $i = 1, \dots, l-1$ and normalized so that $(\tilde{\alpha}_i|\tilde{\alpha}_i) = 1$, $i = 1, \dots, l-1$.

Let $a_i(n)$ ($i = 1, \dots, l$) and $b_i(n)$ ($i = 1, \dots, l-1$) be two set of bosonic operators satisfying

$$\begin{aligned} [a_i(n), a_j(m)] &= \frac{[(\alpha_i|\alpha_j)n]}{n} [n] \delta_{n,-m}, \\ [b_i(n), b_j(m)] &= \frac{[(\alpha_i|\alpha_j)n]}{n} [n] \delta_{n,-m}. \end{aligned}$$

Then level one realization is defined on the Fock space

$$V = \text{Sym}(a_i(-n)'s, b_i(-n)'s) \otimes C[P] \otimes C[\tilde{Q}].$$

The action of the Z -operators are defined by

$$\begin{aligned} Z_i^\pm(z) &= (U_i^+(zq^{-1/2}) + U_i^-(zq^{1/2})(-1)^{2\partial_i})e^{\pm\alpha_i}z^{\pm\partial_i}, \\ Z_l^\pm(z) &= e^{\pm\alpha_l}z^{\pm\partial_l}. \end{aligned} \quad i = 1, \dots, l-1,$$

where the auxiliary operator $U_i^+(z)$ and $U_i^-(z)$ are defined by:

$$U_i^\pm(z) = \exp\left(\pm \sum_{n \geq 1} \frac{b_i(-n)}{[n]} z^n\right) \exp\left(\mp \sum_{n \geq 1} \frac{b_i(-n)}{[n]} z^{-n}\right) e^{\pm\tilde{\alpha}_i} z^{\pm\partial_i}$$

and the operator $e^{\pm\tilde{\alpha}_i}$ acts on $C[A_{l-1}]$ (the group algebra generated by the short roots) by

$$e^{\tilde{\alpha}_i} \cdot e^{\tilde{\beta}} = \epsilon(\alpha_i, \beta) e^{\tilde{\alpha}_i + \tilde{\beta}}.$$

where the function $\epsilon(\alpha_i, \alpha_j)$ is given by

$$\epsilon(\alpha_i, \alpha_j) = \begin{cases} -1, & i = j \\ 1, & i > j \\ (-1)^{a_{ij}}, & i < j \end{cases}$$

To construct the quantum affine algebra $U_q(F_4^{(1)})$, we let $b_i(n)$ ($i = 1, 2$) be the auxiliary bosonic operators satisfying the same relations as the first two Heisenberg generators associated to $U_q(F_4^{(1)})$. Now the short roots form a sublattice isomorphic to that of A_2 . The representation space is defined as

$$V = \text{Sym}(a_i(-n)'s, b_j(-n)'s) \otimes C[P] \otimes C[A_2].$$

where $i = 1, 2$ and $j = 1, 2$.

Similar to the case of $C_n^{(1)}$ we have the following result. Note that we do not need to adjust signs by group algebra.

Theorem 3. 6. *For $i = 1, \dots, 4$ and $j = 1, 2$, The following action defines a level one representation of $U_q(F_4^{(1)})$*

$$\begin{aligned} Z_i^\pm(z) &= (U_i^+(zq^{-1/2}) + U_i^-(zq^{1/2})(-1)^{2\partial_i})e^{\pm\alpha_i}z^{\pm\partial_i}, \\ Z_j^\pm(z) &= e^{\pm\alpha_j}z^{\pm\partial_j}. \end{aligned} \quad i = 1, 2, \quad i = 3, 4,$$

where the auxiliary operator $U_i^+(z)$ and $U_i^-(z)$ are defined by the same expressions as in the case of $C_n^{(1)}$.

It is clear that the operators $U_i^\pm(z)e^{\mp b_i}z^{\mp b_i(0)}$ satisfy the relations of quantum Z -algebras associated with the the same invariant form (|). Thus they are some auxiliary quantum Z -operators in the representation. This phenomena appear in

almost all (vertex operator) reducible modules. To get irreducible modules one can use quantum screening operators [M].

We conclude this subsection by a level $-1/2$ representation of $U_q(C_l^{(1)})$ [JKM1].

Let $a_i(m)$ and $b_i(m)$ ($i = 1, \dots, l; m, n \in \mathbb{Z}$) be the operators satisfying the following relations:

$$\begin{aligned} [a_i(m), a_j(n)] &= \delta_{m+n,0} \frac{[m(\alpha_i|\alpha_j)][-m/2]}{m}, \\ [b_i(m), b_j(n)] &= m\delta_{ij}\delta_{m+n,0} \\ [a_i(m), b_j(n)] &= 0. \end{aligned}$$

The module $\mathcal{F}_{\alpha,\beta} = \mathcal{F}_{\alpha}^1 \otimes \mathcal{F}_{\beta}^2$ is built from the Heisenburg algebra by the defining relations

$$\begin{aligned} a_i(m)|\alpha, \beta\rangle &= 0 \quad (m > 0), \quad b_i(m)|\alpha, \beta\rangle = 0 \quad (m > 0), \\ a_i(0)|\alpha, \beta\rangle &= (\alpha_i|\alpha)|\alpha, \beta\rangle, \quad b_i(0)|\alpha, \beta\rangle = (2\varepsilon_i|\beta)|\alpha, \beta\rangle, \end{aligned}$$

where $|\alpha, \beta\rangle = |\alpha\rangle \otimes |\beta\rangle$ ($\alpha \in P + \frac{\mathbb{Z}}{2}\lambda_l, \beta \in P$) is the vacuum vector. The grading operator d is defined by

$$d|\alpha, \beta\rangle = ((\alpha|\alpha) - (\beta|\beta - \lambda_n))|\alpha, \beta\rangle.$$

We set the Fock space

$$\tilde{\mathcal{F}} := \bigoplus_{\alpha \in P + \frac{1}{2}\mathbb{Z}\lambda_l, \beta \in P} \mathcal{F}_{\alpha,\beta}.$$

Let e^{a_i} and e^{b_i} be operators on $\tilde{\mathcal{F}}$ given by:

$$e^{a_i}|\alpha, \beta\rangle = |\alpha + \alpha_i, \beta\rangle, \quad e^{b_i}|\alpha, \beta\rangle = |\alpha, \beta + \varepsilon_i\rangle.$$

Let $:$ be the usual bosonic normal ordering defined by

$$\begin{aligned} : a_i(m)a_j(n) : &:= a_i(m)a_j(n) \quad (m \leq n), \quad a_j(n)a_i(m) \quad (m > n), \\ : e^a a_i(0) : &:= a_i(0)e^a := e^a a_i(0), \\ : e^b b_i(0) : &:= b_i(0)e^b := e^b b_i(0). \end{aligned}$$

and similar normal products for the $b_i(m)$'s.

Let $\partial = \partial_{q^{1/2}}$ be the q -difference operator:

$$\partial_{q^{1/2}}(f(z)) = \frac{f(q^{1/2}z) - f(q^{-1/2}z)}{(q^{1/2} - q^{-1/2})z}$$

We introduce the following operator:

$$U_i^{\pm}(z) = \exp(\pm \sum_{k=1}^{\infty} \frac{b_i(-k)}{k} z^k) \exp(\mp \sum_{k=1}^{\infty} \frac{b_i(k)}{k} z^{-k}) e^{\pm b_i} z^{\pm b_i(0)}.$$

Proposition 3.7. *The Fock space $\tilde{\mathcal{F}}$ is a U_q -module of level $-\frac{1}{2}$ under the action defined by $\gamma \mapsto q^{-\frac{1}{2}}$, $K_i \mapsto q^{a_i(0)}$, $a_{im} \mapsto a_i(m)$, $q^d \mapsto q^d$, and*

$$\begin{aligned} Z_i^+(z) &\mapsto \partial U_i^+(z) U_{i+1}^-(z) e^{\alpha_i} z^{-2a_i(0)}, & i = 1, \dots, n-1 \\ Z_i^-(z) &\mapsto U_i^-(z) \partial U_{i+1}^+(z) e^{-\alpha_i} z^{2a_i(0)}, & i = 1, \dots, n-1 \\ Z_l^+(z) &\mapsto \left(\frac{1}{q^{\frac{1}{2}} + q^{-\frac{1}{2}}} : U_l^+(z) \partial^2 U_l^+(z) : - : \partial U_l^+(q^{-\frac{1}{2}} z) \partial U_l^+(q^{-\frac{1}{2}} z) : \right) e^{\alpha_l} z^{-2a_l(0)} \\ Z_n^-(z) &\mapsto \frac{1}{q^{\frac{1}{2}} + q^{-\frac{1}{2}}} : U_l^-(q^{\frac{1}{2}} z) U_l^-(q^{-\frac{1}{2}} z) : e^{-\alpha_l} z^{2a_l(0)}. \end{aligned}$$

3.7. Level one realization of $U_q(G_2^{(1)})$ [J3].

We can realize level one $U_q(G_2^{(1)})$ by introducing two sets of auxiliary bosonic operators $b_2(n)$ and $c_2(n)$ ($n \in \mathbb{Z}$) with the defining relations:

$$\begin{aligned} [b_2(n), b_2(m)] &= -\frac{[\frac{2}{3}n]}{n} [n] \delta_{n,-m} \\ [c_2(n), c_2(m)] &= \frac{[\frac{2}{3}n]}{n} [\frac{5}{3}n] \delta_{n,-m} \\ [a_i(n), b_2(n)] &= [a_i(n), c_2(n)] = 0 \end{aligned}$$

Let us enlarge the root lattice by adding an auxiliary root β_2 with $(\beta_2|\beta_2) = -2/3$. The actions of $a_i(m), b_2(m)$ are defined similarly as previous sections. We often identify $a_i(0)$ with ∂_{α_i} .

Proposition 3.8. *On the Fock space*

$$V = \text{Sym}(a_i(-n), b_2(-n), c_2(-n)) \otimes C[P + \mathbb{Z}\beta_2]$$

the correspondence

$$\begin{aligned} Z_1^\pm(z) &\mapsto e^{\pm\alpha_1} z^{\pm\partial_{\alpha_1}} \\ Z_2^\pm(z) &\mapsto e^{\pm\alpha_2} z^{\pm\partial_{\alpha_2}} \frac{U^\pm(z)}{q - q^{-1}} (B_+(zq^{\mp 5/2} z) W_+(q^{\mp 3/2} z)^{\pm 1} \\ &\quad - B_-(zq^{\pm 5/2} z) W_-(q^{\pm 3/2} z)^{\pm 1}) \end{aligned}$$

gives a level one $U_q(G_2^{(1)})$ -module and where

$$\begin{aligned} U^\pm(z) &= \exp(\pm \sum_{m=1}^{\infty} \frac{q^{\mp m/2} b_2(-m)}{[m]} z^m) \exp(\mp \sum_{m=1}^{\infty} \frac{q^{\mp m/2} b_2(m)}{[m]} z^{-m}) e^{\pm\beta_2} z^{\pm b_2(0)} \\ B_\pm(z) &= \exp(\pm(q - q^{-1}) \sum_{m=1}^{\infty} \frac{[m/3]}{[2m/3]} b_2(\pm) z^{\mp m}) q^{\mp b_2(0)/2} \\ W_\pm(z) &= \exp(\pm(q - q^{-1}) \sum_{m=1}^{\infty} \frac{[m/3]}{[2m/3]} c_2(\pm) z^{\mp m}). \end{aligned}$$

The proof is obtained by a careful analysis of the Serre relations [J3].

In this paper we restricted ourselves to the untwisted types or type one. We can similarly introduce the quantum Z-algebras for the twisted types using [J4]. Then all the level one modules of twisted types [J1] can be realized by the method discussed in this paper.

REFERENCES

- [ABH] A. Abada, A.H. Bougourzi, and El. Gradechi, *Deformation of Wakimoto construction*, Mod. Phys. Lett A **8** (1993), 715-724.
- [AOS] H. Awata, S. Odake, and J. Shiraishi, *Free boson representation of $U_q(\hat{\mathfrak{sl}}_N)$* , Commun. Math. Phys. (1994), 61-83.
- [Be] J. Beck, *Braid group action and quantum affine algebras*, Commun. Math. Phys. **165** (1994), 555-568.
- [Br] D. Bernard, *Vertex operator representations of quantum affine algebras $U_q(B^{(1)})$* , Lett. Math. Phys. **17** (1989), 239-245.
- [Bo] A.H. Bougourzi, *Uniqueness of the bosonization of the $U_q(\mathfrak{su}(2)_k)$ quantum current algebra*, Nucl. Phys. **B404** (1993), 457-482.
- [BV] A.H. Bougourzi and H. Vinet, *A quantum analogue of the $\mathcal{Z}$ algebra*, J. Math. Phys. **37** (1996), 3548-3567.
- [DFg] J. Ding and B. Feigin, *Quantum current operators. II. Difference equations of quantum current operators and quantum parafermion construction*, Publ. Res. Inst. Math. Sci. **33** (1997), 285-300.
- [DFr] J. Ding and I.B. Frenkel, *Isomorphisms of two constructions of quantum affine algebras*, Comm. Math. Phys. **156** (1993), 277-300.
- [DW] J. Ding and T. Miwa, *Quantum current operators. I. Zeros and poles of quantum current operators and the condition of quantum integrability*, Publ. Res. Inst. Math. Sci. **33** (1997), 277-284.
- [D] V.G. Drinfeld, *New realization of Yangians and quantum affine algebras*, Sov. Math. Dokl. **36**, 212-216.
- [FJ] I.B. Frenkel and N. Jing, *Vertex representations of quantum affine algebras*, Proc. Nat'l. Acad. USA **85** (1988), 9373-9377.
- [FLM] I. Frenkel, J. Lepowsky, and A. Meurman, *Vertex operator algebras and the Monster*, Academic Press, New York, 1988.
- [J1] N. Jing, *Twisted vertex representations of quantum affine algebras*, Invent. Math. **102** (1990), 663-690.
- [J2] N. Jing, *Higher level representations of the quantum affine algebra $U_q(\hat{\mathfrak{sl}}_2)$* , Jour. Alg. **182** (1996), 448-468.
- [J3] N. Jing, *Level one representations of $U_q(G_2^{(1)})$* , Proc. Amer. Math. Soc., to appear.
- [J4] N. Jing, *On Drinfeld realization of quantum affine algebras*, Proc. Lie algebras and the Monster, W. de Gruyter, Berlin, to appear, q-alg/9610035.
- [JM] N. Jing and K.C. Misra, *Vertex operators of level one $U_q(B_n^{(1)})$ -modules*, Lett. Math. Phys. **36** (1996), 127-143.
- [JKM1] N. Jing, Y. Koyama, and K.C. Misra, *Bosonic realizations of $U_q(C_n^{(1)})$* , Jour. Algebra **200** (1998), 155-172..
- [JKM2] N. Jing, Y. Koyama, and K.C. Misra, *Level one realizations of quantum affine algebras $U_q(C_n^{(1)})$* , submitted (QA/9802123) (1998).
- [I] M. Idzumi, *Level two irreducible representations of $U_q(\hat{\mathfrak{sl}}_2)$* , Int. J. Mod. Phys. A **9** (1994), 4449-4484.
- [K] V.G. Kac, *Infinite dimensional Lie algebras*, 3rd ed., Cambridge University Press, 1990.
- [KMPY] M. Kashiwara, T. Miwa, J.-U.H. Peterson, and C.M Yung, *Perfect crystals and q -deformed Fock spaces*, Selecta Mathematica **2** (1996), 415-499.
- [LP1] J. Lepowsky and M. Primc, *Standard modules for type one affine Lie algebras*, Number Theory, Lect. Notes in Math., vol. 1052, Springer-Verlag, New York, 1984, pp. 194-251.
- [LP2] J. Lepowsky and M. Primc, *Structure of the standard modules for the affine Lie algebra $A_1^{(1)}$* , Contemporary Math., vol. 46, Amer. Math. Soc., Providence, RI, 1985.
- [LW1] J. Lepowsky and R. L. Wilson, *A new family of algebras underlying the Rogers-Ramanujan identities and generalizations*, Proc. Nat'l. Acad. Sci. USA **78** (1981), 7254-7258.
- [LW2] J. Lepowsky and R. L. Wilson, *The structure of standard modules, I: Universal algebras and the Rogers-Ramanujan identities*, Invent. Math. **77** (1984), 199-290.
- [LW3] J. Lepowsky and R. L. Wilson, *The structure of standard modules, II: The case $A_1^{(1)}$* , Invent. Math. **77** (1985), 417-442.

- [M] A. Matsuo, *A q -deformation of Wakimoto modules, primary fields and screening operators*, Comm. Math. Phys. **160** (1994), 33–48.
- [P] M. Primc, *Standard representations of $A_n^{(1)}$* , Infinite-dimensional Lie algebras and groups, World Sci. Singapore, Singapore, 1989, pp. 273–284.
- [S] J. Shiraishi, *Free boson representation of $U_q(\hat{sl}_2)$* , Phys. Lett. A **171** (1992), 243–248.